

THE APPLICATION OF SATELLITE DATA TO MONITOR THE ENCROACHMENT OF AGRICULTURE ON FOREST RESERVE IN THE PHU LUANG WILDLIFE SANCTUARY, LOEI PROVINCE, NE OF THAILAND

Rasamee SUWANWERAKAMTORN¹, Pataraporn PIMDEE¹,
Sirorat MONGKOLSAWAT¹, and Nattaphol SRITOOMKAEW²
Assist. Prof., ¹Center of Geoinformatics for the Development of Northeast Thailand,
Khon Kaen University, Computer Center Building, Khon Kaen, Thailand, 40002.
E-mail: rasamee@kku.ac.th

² Phu Luang Wildlife Sanctuary, Phu Luang, Loei, Thailand, 42230.

KEY WORDS: Landuse Change, Encroachment of Agriculture, Forest Reserve, Phu Luang Wildlife Sanctuary

ABSTRACT: Control and preventive measures are enforced rigidly to minimize the encroachment on forest reserves in the Phu Luang Wildlife Sanctuary (PWLS), a forest area set aside as a wildlife habitat in the North of Thailand, near the Laos border. However, even with these measures in place the forest area is still continuously encroached upon due to a population increase. The purpose of this study was to monitor the encroachment of agriculture on forest reserve in the PWLS. In order to achieve this, remote sensing technology, which is ideal for monitoring the forest change, coupled with multi-dates satellite data were used to meet the objective. Four dates of LANDSAT data acquired in 1994, 1997, 2001, 2005 and two dates of THEOS data acquired in 2010 were used to produce multi-temporal landuse maps in the study area by visual interpretation. These maps were then overlaid using GIS technology to analyze the forest area that had been converted to agricultural land. The result showed that the forest had been gradually depleted by 6,247 ha (6.96%), from 61,069 ha in 1994 to 54,822 ha in 2010. Agriculture had increased by 5,879 ha (6.44%), from 27, 107 ha in 1994 to 32,986 ha in 2010.

1. INTRODUCTION

The Phu Luang Wildlife Sanctuary is one of the 57 wildlife sanctuaries of Thailand. As a sanctuary it is one of the best ways to maintain forest areas as a wildlife habitat. Control and preventive measures are enforced rigidly to minimize the encroachment on forest reserves. Phu Luang Wildlife Sanctuary is considered to be one of the largest sources of flora and fauna in Thailand with a diverse ecosystem and abundant forestry resources that makes it unique in the area (Mongkolsawat, C. et al., 2005; Wildlife Conservation Division and Khon Kaen University, 2000). However, even with the preventive measures in place at the wildlife sanctuary the forest area is still being encroached upon and converted to agricultural land using slash-and-burn techniques. To enhance the efficiency of the protection, regeneration and utilization of forest resources, information about these changes is required. To achieve this, remote sensing technology, which is well suited to this purpose, will be used in conjunction with the high resolution satellite data. The resulting updated and multi-temporal information can be employed to detect the changes. This information can be used for wildlife habitat protection, planning and long-term policy setting.

2. OBJECTIVE

The purpose of this study is to monitor the encroachment of agriculture on a forest reserve with the use of multi-temporal satellite data.

3. THE STUDY AREA

The study area, PLWS is located in the Phu Luang District in the southern part of the Loei Province, NE of Thailand (Figure 1). Its coverage area is approximately 897 square kilometers (89,697 ha). It lies between latitude 17 degrees 3 minutes and 17 degrees 24 minutes North and longitude 101 degrees 16 minutes to 101 degrees 21 minutes East. It is a mountainous area with an escarpment in the East. The height of its summit is about 1,600 m.msl.




Figure 1 Study area

It is characterized by a number of hills with a thick sequence of Mesozoic rock of Phu Kradung, Soa Khua, Pha Dua, Phu Phan and Phra Wihan Formations. The areas support two main forest types: evergreen and deciduous forests. The mean annual rainfall of the area varies from 1,200 – 1,400 mm, with over 60% of the annual rainfall falling in August and September (Mongkolsawat, C. et al., 2005).

4. METHODOLOGY

Details of the methodology are described in the following steps (Figure 2).

4.1 Data sources

4.1.1 Satellite data: Four Landsat data acquired in 1994, 1998, 2001 and 2005 were obtained from the Regional Centre for Geo-Informatics and Space Technology, Northeast Thailand and two scenes of THEOS data, acquired on March 1st and 6th, 2010, from the Geo-Informatics and Space Technology Development Agency (GISTDA) were used to produce multi-temporal landuse maps.

4.1.2 Topographic maps: Topographic maps of the Royal Thai Survey Department at the scale of 1:50,000 were used for geo-referencing and supplementary information.

4.1.3 Existing Landuse maps: Landuse maps from 1998 and 2005 from the Regional Centre for Geo-Informatics and Space Technology, Northeast Thailand were used for reference.

4.2 Data analysis

4.2.1 Pre-processing: Two scenes of THEOS data were mosaiced into one image which was geo-referenced using the ground control points obtained from the topographic map at scale of 1:50,000. A nearest neighbour interpolation algorithm was performed. The four dates of Landsat TM data were co-registered to the geo-coded THEOS image.

4.2.2 Image interpretation: The colour combination of bands 3, 1, 2 (RGB) of the THEOS data and bands 4, 5, 3 (RGB) of the Landsat TM data were visually interpreted landuse categories. ARCGIS was used to delineate landuse type. A hierarchical interpretative scheme was devised which consisted of the following steps; firstly, directly-observable features such as water bodies water channels, roads and community areas were identified; secondly, stratification of the target area, based on colour, texture and pattern of the imagery was carried out and general features of the area being mapped were noted. Sampling sites for field survey are also identified at this stage; landuse patterns were identified with particular reference to the classification schemes as used by the Land Development Department (Mongkolsawat, C. et al., 2005)

4.2.3 Validation: Field surveys were carried out to increase map accuracy. The field investigation provided site characteristics as related to the image features. The 221 ground sites of different landuse categories were explored to identify cover type and their composition, topography and land types as related to image elements. The obtained map was checked against the ground-truth data and a confusion matrix was applied to validate the obtained results.

4.2.4 Encroachment of Agriculture on Forest Reserve Analysis: A comparison of a pair of landuse maps between the year 1994 and 1998, 1998 and 2001, 2001 and 2005, and 2005 and 2010 was carried out to obtain the landuse change maps. The statistics of the changed class were calculated. The intention of this study was to monitor the change of forest and agricultural classes.

5. RESULT AND DISCUSSIONS

The landuse maps and the acreage of the PLWS for the years 1994, 1998, 2001, 2005, 2010, based on the multi-temporal data of Landsat TM and THEOS are shown in Figure 3 and table 1. The encroachment of agricultural areas on the forest areas in the PLWS had greatly increased from about 30.22% in 1994 to 36.77% in 2010, representing agricultural areas in the PLWS of about 32,986 ha for the year 2010. The total forest areas in the PLWS accounted for 68.08%, 64.60%, 63.27%, 61.89% and 61.12 % for the years 1994, 1998, 2001, 2005 and 2010 respectively. The obtained results indicate that the majority of the forest type was restricted to hill evergreen and dry evergreen forests, with over 70% of the total forest area. To date, in the PLWS the forest cover represents about 61.12%. The depletion in forest areas by type was 1.51%, 2.85% and 1% for hill evergreen, dry evergreen and dry dipterocarp forests respectively (Figure 4 and Table 2). The rate of forest change in the PLWS for the

periods 1994-1998, 1998-2001, 2001-2005 and 2005-2010 accounted for -3.48%, -1.33%, -1.38% and -0.77% respectively, with an overall depletion rate (1994-2010) of 6.96% (Table 2). Significant changes were found during the period 1994-1998 with a progressive decrease in changes after 1998, during which the preventive measures were rigidly enforced. The encroachment on the forest reserves resulted in a substantial expansion of agriculture covering an area of about 5,779 ha (6.44%) from 1994-2010.




Figure 2 The methodology procedure




Figure 3 Landuse maps of PLWS from 1994 to 2010

Table 1 Statistical data of each landuse type in PLWS from 1994 to 2010

Landuse Type	Area of Landuse in Wildlife Sanctuary									
	1994		1998		2001		2005		2010	
	ha	%	ha	%	ha	%	ha	%	ha	%
Forest	61,069	68.08	57,953	64.60	56,755	63.27	55,508	61.89	54,822	61.12
-Hill evergreen forest	25,739	28.70	25,466	28.38	24,984	27.85	24,626	27.45	24,392	27.19
-Pine forest	650	0.72	665	0.74	726	0.81	722	0.80	752	0.84
-Dry or semi-evergreen	21,139	23.56	18,682	20.83	18,707	20.85	18,629	20.77	18,578	20.71
-Mixed deciduous forest	2,725	3.04	2,813	3.14	2,687	3.00	2,725	3.04	2,717	3.03
-Dry dipterocarp forest	2,646	2.95	2,089	2.33	1,799	2.01	1,724	1.93	1,749	1.95
- Bamboo forest	1,083	1.21	1,004	1.12	1,192	1.33	1,223	1.36	1,252	1.40
- Forest Plantation	7,087	7.90	7,234	8.06	6,660	7.42	5,859	6.54	5,382	6.00
Range land	1,077	1.20	1,104	1.23	1,287	1.43	1,314	1.46	1,382	1.54
Outcrops	221	0.25	221	0.25	221	0.25	222	0.25	221	0.25
Agriculture	27,107	30.22	30,193	33.66	31,166	34.75	32,373	36.09	32,986	36.77
Community	208	0.23	211	0.24	234	0.26	246	0.27	252	0.28
Water body	15	0.02	15	0.02	34	0.04	34	0.04	34	0.04
Total	89,697	100.00	89,697	100.00	89,697	100.00	89,697	100.00	89,697	100.00

Source : Data in year 1998 and 2005 modified from Mongkolsawat, C. et al., 2005.




Figure 4 Landuse change in PLWS 1994-2010

Table 2 Landuse change in PLWS from 1994 to 2010

Landuse Type	Area of Landuse change in Wildlife Sanctuary									
	1994-1998		1998-2001		2001-2005		2005-2010		1994-2010	
	ha	%	ha	%	ha	%	ha	%	ha	%
Forest	-3,116	-3.48	-1198.00	-1.33	-1247.00	-1.38	-686.00	-0.77	-6247.00	-6.96
-Hill evergreen forest	-273	-0.32	-482.00	-0.53	-358.00	-0.40	-234.00	-0.26	-1347.00	-1.51
-Pine forest	15	0.02	61.00	0.07	-4.00	-0.01	30.00	0.04	102.00	0.12
-Dry or semi-evergreen	-2,457	-2.73	25.00	0.02	-78.00	-0.08	-51.00	-0.06	-2561.00	-2.85
-Mixed deciduous forest	88	0.10	-126.00	-0.14	38.00	0.04	-8.00	-0.01	-8.00	-0.01
-Dry dipterocarp forest	-557	-0.62	-290.00	-0.32	-75.00	-0.08	25.00	0.02	-897.00	-1.00
- Bamboo forest	-79	-0.09	188.00	0.21	31.00	0.03	29.00	0.04	169.00	0.19
- Forest Plantation	147	0.16	-574.00	-0.64	-801.00	-0.88	-477.00	-0.54	-1705.00	-1.90
Range land	27	0.03	183.00	0.20	27.00	0.03	68.00	0.08	305.00	0.34
Outcrops	-	-	-	-	-	-	-	-	-	-
Agriculture	3,086	3.44	973.00	1.09	1207.00	1.34	613.00	0.68	5879.00	6.55
Community	3	0.01	23.00	0.02	12.00	0.01	6.00	0.01	44.00	0.05
Water body	-	-	19.00	0.02	-	-	-	-	19.00	0.02

6. CONCLUSIONS

Using multi-temporal satellite data the periodical monitoring and management of the PLWS can be effectively carried out. A significant decrease in the rate in which forest encroached upon was seen over the past five years. Continuing control and preventive measures to protect the forest should be made, employing spatial information acquired from periodical satellite data.

7. BIBLIOGRAPHY

- Gibson, P.J., and Power, C.H., 2000. *Introductory Remote Sensing Digital Processing and Applications*. London : Routledge.
- Gu, Y., Brown, J.F., Verdin, J.P. and Wardlow, B., 2007. A five-year analysis of MODIS NDVI and NDWI for grassland drought assessment over the central Great Plains of the United States [Electronic version]. *Geophysical Research Letters*, 34, L 06407.
- Huete, A.R., Didan, K., Miura, T., Rodriguez, E.P., Gao, X. and Ferreira, L.G., 2002. Overview of the radiometric and biophysical performance of the MODIS vegetation [Electronic version], *Remote Sensing of Environment*, 83, 195-213.
- Khon Kaen University and Phu Khieo Wildlife Sanctuary., 1995. *Integrated Information System for the Management of Phu Khieo Wildlife Sanctuary*. Final report submitted to CIDA-Thailand.
- Mongkolsawat, C., Putklang, W., and Suwanwerakamtorn, R., 2005. Forest Change Detection Using Multi-temporal Remotely Sensed Data in Phu Luang Wildlife Sanctuary, Northeast, Thailand. *Proceeding of the 26th Asian Conference on Remote Sensing*, November 7-11, 2005, Hanoi, Vietnam.
- Thailand Institute of Scientific and Technological Research., 1992. *General Information of Master Plan Phu Phan Nation Park*, Kalasin. Sakon Nakhon Province, Bangkok.
- Volcani, A., Karnieli, A. and Svoray, T. ,2005. The use of remote sensing and GIS for spatio-temporal analysis of the physiological state of a semi-arid forest with respect to drought years [Electronic version]. *Forest Ecology and Management*. 215,239-250.
- Weier, J., and Hering, D., 1999. *Measuring Vegetation (NDVI&EVI)*. Retrieved December 3, 2005, from: http://earthobservatory.nasa.gov/Library/Measuring_Vegetation/printal.php
- Wildlife Conservation Division and Khon Kaen University., 2000. *Application of Remote Sensing and GIS for Monitoring Forest Land Use Change in Phu Luang Wildlife Sanctuary*. Final Report submitted to the Royal Forest Department.